

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 17/08, A61K 31/70		A1	(11) International Publication Number: WO 96/19489 (43) International Publication Date: 27 June 1996 (27.06.96)
(21) International Application Number: PCT/IE95/00065 (22) International Filing Date: 19 December 1995 (19.12.95) (30) Priority Data: 940973 19 December 1994 (19.12.94) IE (71) Applicant (for all designated States except US): RUSSINSKY LIMITED [IE/IE]; 90 South Mall, Cork (IE). (72) Inventors; and (75) Inventors/Applicants (for US only): SCHICKANEDER, Helmut [DE/IE]; "Dunhamon", South Douglas Road, Cork (IE). NIKOLOPOULOS, Aggelos [DE/IE]; Apartment No. 3 Sydenham, Off Wellington Road, Cork (IE). KELLY, Declan [IE/IE]; Ringacoltie, Rushbrooke, Cobh, County Cork (IE). (74) Agents: O'CONNOR, Donal, H. et al.; Cruickshank & Co., 1 Holles Street, Dublin 2 (IE).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DE (Utility model), DK, DK (Utility model), EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: COMPOUNDS (57) Abstract Antibiotic and mucolytic salts of roxithromycin, clarithromycin and azithromycin are described. Particularly described is roxithromycin stinoprate which was found to be more potent than erythromycin stinoprate.			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

"Compounds"Introduction


The present invention relates to new antibiotic and mucolytic salts of roxithromycin, clarithromycin and azithromycin.

- 5 The exploitation of the therapeutical properties of thiolic compounds in combination with the properties of antibiotics has been attempted as described in EP 0,057,489A. However, it has been found that acetyl-
10 cysteine and the derivatives thereof are relatively unstable and especially sensitive to oxygen, sunlight, humidity and heat.

The invention is directed towards providing antibiotic and mucolytic salts.

Statements of Invention

- 15 The derivatives according to the present invention have the following general formula :



wherein R is a radical selected from :

Roxithromycin, Clarithromycin and Azithromycin.


- 20 It has been surprisingly found that the compounds of the present invention are very stable, have very low toxicity and can be therapeutically used once daily.

- 2 -


These new derivatives of the below mentioned macrolides have the further advantages of :-

- 5 (a) better oral absorption;
(b) faster and superior concentration;
(c) slow elimination; and/or
(d) reduced dosage : only 300 - 500 mg per day.

Roxithromycin has the structure




Clarithromycin has the structure



- 3 -

Azithromycin has the structure



The compounds according to the invention are white microcrystalline powders.

5 Their use is foreseen in all pharmaceutical forms and the compounds may be provided in any suitable pharmaceutical composition including : capsules; solutions; injectable preparations; aerosols; effervescent tablets; powders; creams; and suspensions. The pharmaceutical compositions will typically contain suitable excipients and/or vehicles
10 which are conventionally used in galenical pharmacy.

The method for the preparation of the new salts comprises reacting roxithromycin, clarithromycin or azithromycin base with N-acetylcystein in a stoichiometrical ratio or preferably with a slight excess of the antibiotic nucleus.
15 Most preferably the reaction is carried out in an organic solvent, at a temperature of between 20 and 40°C and in the presence of water, preferably in an amount of not greater than 20%. Alternatively, the reactions may be carried out in a suspension of water at a temperature of 20 and 40°C,
20 (with N-acetylcystein in a stoichiometrical ratio or in

- 4 -

the presence of a slight excess of the antibiotic nucleus), and after building the salts, water is distilled off under very mild conditions (low vacuum, low temperature).

5 **Example**

Salt of roxithromycin with N-acetylcystein

1800 g of Roxithromycin and 350.9 g of N-acetylcystein are homogenised under inert conditions (nitrogen) for 60 minutes at 15-20°C. To this mixture 720 ml of deionised
10 water is added at atmospheric pressure and further homogenised for one hour at 15-20°C. The product is dried under vacuum and milled, if necessary. A yield of 79.8% (1716 g) is obtained.

15 The Infra Red spectrum of the compound is plotted in Fig. 1.

The salts of clarithromycin and azithromycin are produced in a similar manner to that described in the above example.

MICROBIOLOGICAL ASSAY OF ROXITHROMYCIN STINOPRATE

20 References: USP 23 <81> Antibiotics-Microbial Assays, page 1690-1696 Code of Federal Regulations Title 21 :436.100 - 436.106

Test Organism:

25 Staphylococcus aureus ATCC 29737
The test organism was maintained through periodic inoculations on agar slants containing USP 23 Medium No. 1. The slants are incubated at 32-35°C for 24 hours, and

- 5 -

stored under refrigeration.

Assay Receptacles:

5 Sterile plastic petri-dishes (ca. 20 x 100mm) with covers were used as assay plates. Assay cylinders were manufactured from stainless steel (o.d. 8mm \pm 0.1mm, i.d. 6mm \pm 0.1mm, length 10mm \pm 0.1mm).

Inoculum Preparation:

10 Using 5ml of sterile USP Saline T.S. the growth from an agar slant of S.aureus was washed and made up to 50ml with sterile saline solution. This stock suspension was diluted with sterile saline so that the transmittance, at 580nm, was 25% against
15 saline as the blank.

1ml of this solution (i.e. giving 25% transmission) was added to each 100ml of culture media (USP 23 Medium No. 1 (Oxoid Antibiotic Medium No. 1)).

20 Culture Medium:

Oxoid Antibiotic Medium No. 1.

Solutions: Saline Solution (sterile)

Sodium chloride	0.9g
Purified Water	100ml

25 Sterilised at 121°C for 20 minutes.

Buffer Solution No. 3 (USP 23)

(0.1M potassium phosphate buffer pH 8.0)	
Dibasic potassium phosphate	16.73g
Monobasic potassium phosphate	0.523g

- 6 -

Purified Water 1000ml

Adjusted with 10N potassium hydroxide to give a pH 7.9-8.1. Sterilised at 121°C for 20 minutes.

5 Standards: Erythromycin Stinoprate Standards:

10 About 100mg of Erythromycin Stinoprate standard, accurately weighed, was added to a 100ml volumetric flask. 10ml of methanol was added to dissolve the Erythromycin Stinoprate. This solution contains 10mg/ml of Erythromycin Stinoprate. A 1:10 dilution of this solution was then prepared with sterile buffer solution to obtain a 1000 µg/ml Erythromycin Stinoprate solution.

15

From this stock solution the following dilutions were prepared:

1.56 µg/ml (S_5); 1.25 µg/ml (S_4); 1.0 µg/ml (S_3); 0.8 µg/ml (S_2); and 0.64 µg/ml (S_1) (using sterile buffer solution for dilutions)

20

Test Sample (Roxithromycin Stinoprate):

25 About 100mg of Roxithromycin embonate test substance, accurately weighed, was added to a 100ml volumetric flask. 10ml of methanol was added for dissolution and this in turn was diluted with sterile buffer solution to obtain a 1000 µg/ml Roxithromycin Stinoprate solution.

- 7 -

From this stock solution a solution was prepared containing 1.0 µg/ml Roxithromycin Stinoprate, using sterile buffer solution for dilution purposes (U₃).

Method:

Cylinder-Plate Method USP 23 <81>

Approximately 20ml of sterilised Oxoid Antibiotic Medium No. 1 was placed in each of 22 sterile petri-dishes, and allowed to harden. Using the inoculum described above 5.0ml of seed layer inoculum were added to each plate except two plates which were reserved as "negative controls". Six (6) stainless-steel assay cylinders were dropped on the inoculated surfaces of 18 of the plates from a height of 12mm, with even spacing on a radius of 28mm. The remaining two inoculated plates were retained as positive controls.

Using a 1-level assay with standard curve, alternate cylinders on each of three plates were filled with the 1.0 µg/ml solution of Erythromycin Stinoprate (S₃), and each of the remaining nine cylinders were filled with one of the four other dilutions of the Standard (S₁ - S₅). This process was repeated for the other three dilutions of the standard.

The 1.0 µg/ml solution of Erythromycin Stinoprate (S₃) was filled into alternate cylinders on each of three plates and

- 8 -

the remaining nine cylinders were filled with the Test Sample (U_3).

Incubation: 24 hours at 32-35°C

Estimation of Potency:

5 The potency of the Roxithromycin
Stinoprate test substance was compared
to the standard curve obtained for the
Erythromycin Stinoprate standard
10 dilutions. This was calculated as a
percentage of the potency of the 1.0
 $\mu\text{g/ml}$ Erythromycin Stinoprate standard
dilution.


Results: Roxithromycin Stinoprate

15 Mean potency (\bar{x}) = 114% (of Erythromycin
Stinoprate).
Standard deviation (s.d.) = 2.94%.
Coefficient of variation (c.v.) = 2.56%.

20 The invention is not limited to the embodiments
hereinbefore described which may be varied in detail.

CLAIMS


1. A compound of the formula



wherein R is a radical selected from :

Roxithromycin, Clarithromycin and Azithromycin.

- 5 2. N-Acetylcystein-Roxithromycin Salt.
3. N-Acetylcystein-Clarithromycin Salt.
4. N-Acetylcystein-Azithromycin Salt.
5. A compound substantially as hereinbefore described with reference to the examples.
- 10 6. A pharmaceutical composition comprising a compound as claimed in any preceding claim together with at least one pharmaceutically acceptable excipient and/or carrier.
7. A process for preparing a compound of the formula



- 10 -

wherein R is a radical selected from :

Roxithromycin, Clarithromycin and Azithromycin.

comprising the step of reacting N-Acetylcystein
with roxithromycin, clarithromycin or azithromycin
base.

5

8. A process as claimed in claim 7 wherein the
reaction is carried out in an organic solvent.

9. A process as claimed in claim 7 or 8 wherein the
reaction is carried out in an aqueous medium.

10 10. A process as claimed in any of claims 7 to 9
wherein the reaction is carried out at a
temperature of from 20°C to 40°C.

11. A process substantially as hereinbefore described
with reference to the Examples.

15 12. A compound whenever prepared by a process as
claimed in any of claims 7 to 11.




Fig. 1

INTERNATIONAL SEARCH REPORT

Internatic Application No

PCT/IE 95/00065

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07H17/08 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 057 489 (REFARMED RECH PHARM MED) 11 August 1982 cited in the application see the whole document ---	1-12
Y	INT. J. CLIN. PHARMACOL. THER. TOXICOL., vol. 26, 1988, pages 444-7, XP002002013 M. DE BERNARDI ET AL.: "Human pharmacokinetics of erythromycin propionate-N-acetylcysteinate" see the whole document --- -/--	1-12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

8 May 1996

Date of mailing of the international search report

30-05-1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Bardili, W

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IE 95/00065

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT.

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EUR. J. RESPIR., vol. 1, 1988, pages 171-5, XP002002014 O. PAULSEN ET AL.: "No effect of oral N-acetylcysteine on the bioavailability of erythromycin and bacampicillin" see the whole document ---	1-12
Y	ANNAL. PHARMACOTHER., vol. 26, 1992, pages 46-55, XP002002015 N. BAHAL AND M.C. NAHATA: "The new macrolide antibiotics: azithromycin, clarithromycin, dirithromycin and roxithromycin" see the whole document ---	1-12
Y	J. CLIN. MICROBIOL., vol. 5, 1977, pages 58-61, XP002002481 M.F. PARRY AND H. NEU: "Effect of N-Acetylcysteine on Antibiotic Activity and Bacterial Growth In Vitro" see the whole document ---	1-12
A	FR,A,2 534 588 (ROUSSEL-UCLAF) 20 April 1984 see the whole document -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat Application No

PCT/IE 95/00065

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0057489	11-08-82	FR-A- 2502627	01-10-82
		AT-T- 11546	15-02-85
		EP-A,B 0060573	22-09-82
		EP-A,B 0174395	19-03-86
		JP-C- 1665945	29-05-92
		JP-B- 3017840	11-03-91
		JP-A- 57176996	30-10-82
		JP-C- 1594100	14-12-90
		JP-B- 2016320	16-04-90
		JP-A- 57176997	30-10-82
		JP-C- 1593607	14-12-90
		JP-B- 2016758	18-04-90
		JP-A- 60132999	16-07-85
		US-A- 4393053	12-07-83

FR-A-2534588	20-04-84	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.